

LEVEL-TWO SURVEY DATA SHEET



(1) Determine the stream-reach boundary. (2) Near the lower end of the reach (in the deepest portion of the run), collect water samples and analyze using the chemical tests you have available. You may use your collection container to observe watercolor and clarity and to determine water odors. (3) Measure the width-depth, velocity and estimate the water level. (4) If you use a two-pole **kick-net**, collect a minimum of **three** benthic macroinvertebrate samples from the best riffle or runs within your stream reach. Use the table on page five to record information about your collections. (5) Evaluate the physical and habitat conditions; record information about known land use activities. (6) Sketch your reach or submit photographs and add any other comments that you feel are important. Note: A WVDNR Scientific Collection Permit is required for all benthic collections.

Stream name _____ Survey date _____
 Watershed _____ Station code _____
 Latitude _____ Longitude _____ Directions to site _____

Survey completed by _____
 Current weather conditions _____
 Past weather conditions (last 3-days) _____
 Affiliation _____ Email _____
 Mailing _____ Phone number _____
 address _____

WATER CHEMISTRY: Use the spaces below to record the results of your water chemistry analysis; attach additional sheets if necessary.

	Result	units		Result	units		Result	units
Temperature (C/F)			Conductivity			Alkalinity		
Dissolved oxygen			Nitrates			Iron		
pH			Turbidity			Fecal/E-coli		
Additional tests (describe and record results) _____								

PHYSICAL CONDITIONS: Use the check boxes below to describe the conditions that closely resemble those of your stream. The extra lines are provided to write in any additional comments. You may see more than one type of condition; if so, be sure to indicate these on your survey (check all that apply). If multiple conditions are observed, always indicate the most dominant condition. If the condition you observe is not listed, describe it in the comment section.

Water clarity	Watercolor	Water/sediment odor	Surface foam
		Water	Sediment
Clear	None	None	None
Murky	Brown	Fishy	Slight
Milky	Black	Musky	Moderate
Muddy	Orange/red	Rotten egg	High
Other (describe)	Gray/White	Sewage	
	Green	Chemical	

Algae color	Algae abundance	Algae growth habit	Streambed color
Light green	None	Even coating	Brown
Dark green	Scattered	Hairy	Black
Brown	Moderate	Matted	Green
Other (describe)	Heavy	Floating	White/gray
			Orange/red

Estimate and indicate the percentage of your reach that is shaded.

> 80	80-60	60-40	< 40
Excellent	Good	Marginal	Poor

Physical condition comments: _____

LEVEL-TWO SURVEY DATA SHEET

WIDTH AND DEPTH: Record the wetted width and depth of the channel's features (riffles, runs or pools). Choose two or more features to measure. Record the average depth from a minimum of four measurements (one of these should be from the deepest part of the feature). The width should be measured from the widest section of the feature.

- 1. Riffle Wetted width ^(feet) _____ Depth ^(feet) _____
- 2. Run Wetted width ^(feet) _____ Depth ^(feet) _____
- 3. Pool Wetted width ^(feet) _____ Depth ^(feet) _____

DISCHARGE

Determine the discharge by using a flow meter (if available) or other methods such as the **FLOAT** or a **VELOCITY HEAD ROD (VHR)**. Discharge should be measured from a run (area of the channel with fast moving water with no breaks in the surface such as protruding rocks). The more measurements collected the more accurate your discharge results will be. To convert inches into feet, divide by 12. For example, if your depth measurement was 6-inches the result in feet would be 0.5. Indicate the methods chosen to measure the discharge and use the tables to record your results. Use the table on the next page to record your measurements.

Discharge method used

Water Level

Float
 VHR
 Flow meter

Low
 Normal
 High
 Dry

Channel width _____ feet

Use the table to record your velocity data

Distance (ft)	Depth (ft)	Velocity (ft/sec)	VHR (Rise-inches)	Float (sec)	Discharge (cfs)
1					
2					
3					
4					
5					
6					
7					
8					
9					
10					
11					
12					
13					
14					
15					

Average Depth _____ feet

Cross Sectional Area (CSA) _____ ft²
 (CSA = Average Depth x Width)

Discharge = CSA x Velocity
 = _____ x _____
 = _____ cfs (ft³/sec)

If you use a float record your distance below and the number of seconds, it took to travel the distance in the column indicated.

Float distance (feet) _____

Use the table below to determine **VHR velocity** from the rises recorded above. The rises below are in inches.

Rise (R)	Velocity	Rise (R)	Velocity
1/4	1.2	3 1/4	4.2
1/2	1.6	3 1/2	4.3
3/4	2.0	3 3/4	4.5
1	2.3	4	4.6
1 1/4	2.6	4 1/4	4.8
1 1/2	2.8	4 1/2	4.9
1 3/4	3.1	4 3/4	5.0
2	3.3	5	5.2
2 1/4	3.5	5 1/4	5.3
2 1/2	3.7	5 1/2	5.4
2 3/4	3.8	5 3/4	5.5
3	4.0	6	5.7

VHR Velocity = $8 \times \sqrt{R}$, where R is rise in feet.

LEVEL-TWO SURVEY DATA SHEET

CHANNEL PROFILES: Width and depth measurements can be used to create a cross section profile within your reach. Choose a location in your reach across one of the channel types above. Stretch a tape from bank to bank and anchor it at both ends. Move from left to right facing in an upstream direction; measure the distance from the stream bottom to the top of the tape at selected intervals (i.e., every foot). Record your measurements in the table below. The table provides enough spaces for 20 measurements; if more are necessary you can create your own table on a separate piece of paper. Your tape measure will probably not start at zero so make sure to record the actual position of the tape as you measure across the channel.

Width intervals


1	2	3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18	19	20

Depth measurements

1	2	3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18	19	20

PEBBLE COUNT: Collect a minimum of 100-particles from your reach using a Zigzag method, percent habitat method or specific transects (e.g., every 10-meter). If you do not complete a pebble count, **ALWAYS ESTIMATE** streambed composition from the riffles/runs chose for your macroinvertebrate sample collections.

Indicate your method from the choices below.	Size Classes (Intermediate axis in millimeters)						
	Silt/clay < 0.06	Sand 0.06 – 2	Fine Gravel 2 – 24	Coarse Gravel 25 – 64	Cobble 65 – 255	Boulder 256 – 1096	Bedrock > 1096
Zigzag							
% Habitat							
10-m Transects							
Woody Debris Includes sticks, roots, leaves etc.							
Totals							



- (A) Long axis (**Length**)
- (B) Intermediate axis (**Width**)
- (C) Short axis (**Height**)





Pebble counts require two people, one in the stream and one on shore. The person in the stream slowly walks upstream from bank to bank using one of the methods above. After each step, the person reaches down without looking, picks up the first particle touched, and measures the intermediate axis with a ruler. The on-shore partner records the measurement. The process continues until 100 pebbles have been measured or the reach has been walked.

HABITAT CONDITIONS: Score each habitat condition using the scales provided. Add all the scores to determine your overall habitat score and integrity rating. Feel free to describe additional features that you feel are important.

LEVEL-TWO SURVEY DATA SHEET

Sediment deposition	Little or no formation of depositional features; < 20% of the reach affected.					Some increase in depositional features; 20-40% of the reach affected.					Moderate amounts of depositional features; 40-60% of the reach affected.					Heavy amounts of deposition; > 60% of the reach affected.				
	Score	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2

Embeddedness should be evaluated in **RIFLES** prior to or during your macroinvertebrate collections.

Embeddedness																				
	<p>Fine sediments surrounds <10% of the spaces between the gravel, cobble, and boulders.</p> <p>Fine sediment surrounds 10-30% of the spaces between the gravel, cobble, and boulders.</p> <p>Fine sediment surrounds 30-60% of the spaces between the gravel, cobble, and boulders.</p> <p>Fine sediment surrounds > 60% of the spaces between the gravel, cobble, and boulders.</p>																			
Score	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1

The last three conditions are assessed on both sides of the channel. The **LEFT** and **RIGHT** sides are determined by looking downstream.

Bank vegetative protection	> 90% of the banks are covered by natural vegetation; all levels (trees, shrubs and herbs) represented; disruption from grazing, mowing etc. minimal or absent; all plants allowed to grow naturally.					70-90% of the banks covered by natural vegetation; one level of plants may be missing or not well represented; some disruption of vegetation evident; > 50% of the potential plant height remains.					50-70% of the banks covered by natural vegetation; patches of bare soil may be present and closely cropped vegetation is common; < 50% of the potential plant heights remains.					< 50% of the banks covered by natural vegetation; disruption is high; vegetation has been removed or the potential plant heights are greatly reduced.				
	Left	10	9	8	7	6	5	4	3	2	1	10	9	8	7	6	5	4	3	2
Right	10	9	8	7	6	5	4	3	2	1	10	9	8	7	6	5	4	3	2	1
Bank stability	Banks are stable; no evidence of erosion or bank failure; little or no potential for future problems.					Banks are moderately stable; infrequent areas of erosion occur, mostly shown by banks healed over.					Banks are moderately unstable; 60% of the reach has some areas of erosion; high potential for erosion during flooding events.					Banks are unstable; many have eroded areas (bare soils) along straight sections or bends; obvious bank collapse or failure; > 60% of the reach has erosion scars.				
	Left	10	9	8	7	6	5	4	3	2	1	10	9	8	7	6	5	4	3	2
Right	10	9	8	7	6	5	4	3	2	1	10	9	8	7	6	5	4	3	2	1
Riparian buffer width	Mainly undisturbed vegetation > 60 ft; no evidence of human impacts such as parking lots, roadbeds, clear-cuts, mowed areas, crops, lawns etc.					Zone of undisturbed vegetation 40-60 ft; some areas of disturbance evident.					Zone of undisturbed vegetation 20-40 ft; disturbed areas common throughout the reach.					Zone of undisturbed vegetation < 20 ft; disturbed areas common throughout the entire reach.				
	Left	10	9	8	7	6	5	4	3	2	1	10	9	8	7	6	5	4	3	2
Right	10	9	8	7	6	5	4	3	2	1	10	9	8	7	6	5	4	3	2	1
Total score	> 85					85 - 70					69 - 50					< 50				
Integrity rating	Optimal					Suboptimal					Marginal					Poor				

SEDIMENT DEPOSITION MAY CAUSE THE FORMATION OF ISLANDS, POINT BARS (AREAS OF INCREASED DEPOSITION USUALLY AT THE BEGINNING OF A MEANDER THAT INCREASE IN SIZE AS THE CHANNEL IS DIVERTED TOWARD THE OUTER BANK) OR SHOALS, OR RESULT IN THE FILLING OF RUNS AND POOLS. USUALLY, DEPOSITION IS EVIDENT IN AREAS THAT ARE OBSTRUCTED BY NATURAL OR MANMADE DEBRIS AND AREAS WHERE THE STREAM FLOW DECREASES, SUCH AS BENDS.

Habitat comments: _____

LEVEL-TWO SURVEY DATA SHEET

LAND USE: Indicate the land uses that you believe may be having an impact on your stream station. Use the letters **(S)** streamside, **(M)** within 1/4 mile and **(W)** somewhere in the watershed, to indicate the approximate location of the disturbance and the numbers **(1)** slight, **(2)** moderate or **(3)** high, to represent the level of disturbance.

Active Construction		Pastureland		Single-family residences	
Mountaintop mining		Cropland		Sub-urban developments	
Deep mining		Intensive feedlots		Parking lots, strip-malls etc.	
Abandoned mining		Unpaved Roads		Paved Roads	
Logging		Trash dumps		Bridges	
Oil and gas wells		Landfills		Other (describe)	
Recreation (parks, trails etc.)		Industrial areas			

Land use comments: _____ Pipes? Yes No

Describe the types of pipes observed and indicate if there is any discharge from the pipes. Also describe the colors and odors of the discharge.

PHOTOGRAPH and SKETCH YOUR REACH: Use the space below or a separate piece of paper to draw your study reach. Indicate the direction of flow, north, sample locations and important features of the reach. Photographs are an excellent method for tracking changes, especially changes related to the condition of the habitat. Choose a minimum of two permanent locations from which to take your photos. Submit your photos with your survey data sheet.

LEVEL-TWO SURVEY DATA SHEET

BENTHIC MACROINVERTEBRATES

Assess your macroinvertebrate collections by counting and identifying to the family-level if possible. Use the table on the **below** to record your collections data. Although streamside identification is possible at this level, WV Save Our Stream's recommends preserving your samples using a full count or standard sub-sampling procedure in a well-lit and more comfortable setting.

The dot-dash tally method is a convenient way to record your data. Each dot or dash represents one tally.

1 2 3 4 5 6 7 8 9 10
 • •• ••• •••• ••••• •••••• •••••• ••••••• ••••••••

INSECT GROUPS

Patterned stoneflies Taxa <input type="text"/> Total <input type="text"/>	Winter stoneflies Taxa <input type="text"/> Total <input type="text"/>	Roach-like stonefly Total <input type="text"/>
Giant stonefly Total <input type="text"/>	Brown stonefly Total <input type="text"/>	Spiny crawler mayfly Total <input type="text"/>
Square-gilled mayfly Total <input type="text"/>	Minnow mayflies Taxa <input type="text"/> Total <input type="text"/>	Flatheaded mayfly Total <input type="text"/>
Brush-legged mayfly Total <input type="text"/>	Burrowing mayflies Taxa <input type="text"/> Total <input type="text"/>	Net-spinning caddisflies Taxa <input type="text"/> Total <input type="text"/>
Case-building caddisflies Taxa <input type="text"/> Total <input type="text"/>	Free-living caddisfly Total <input type="text"/>	Common netspinner Total <input type="text"/>
Dragonflies Taxa <input type="text"/> Total <input type="text"/>	Damselflies Taxa <input type="text"/> Total <input type="text"/>	Riffle beetle Total <input type="text"/>
Long-toed beetle Total <input type="text"/>	Water penny Total <input type="text"/>	Other beetles (true bugs) Taxa <input type="text"/> Total <input type="text"/>
Hellgrammite/Fishfly Total <input type="text"/>	Alderfly Total <input type="text"/>	Aquatic moth Total <input type="text"/>

Continue recording on the next page.

LEVEL-TWO SURVEY DATA SHEET

Non-biting midge Total <input type="text"/>	Black fly Total <input type="text"/>	Crane fly Total <input type="text"/>
Watersnipe fly Total <input type="text"/>	Dance fly Total <input type="text"/>	Dixid midge Total <input type="text"/>
Net-wing midge Total <input type="text"/>	Horse fly Total <input type="text"/>	Other fly larva Taxa <input type="text"/> Total <input type="text"/>

NON-INSECT GROUPS

Crayfish Total <input type="text"/>	Scud/Sideswimmer Total <input type="text"/>	Aquatic sowbug Total <input type="text"/>
Water mite Total <input type="text"/>	Operculate snails Taxa <input type="text"/> Total <input type="text"/>	Non-operculate snails Taxa <input type="text"/> Total <input type="text"/>
Pea clam Total <input type="text"/>	Asian clam Total <input type="text"/>	Mussel Total <input type="text"/>
Flatworms Total <input type="text"/>	Aquatic worms Total <input type="text"/>	Leeches Total <input type="text"/>

Other aquatic invertebrates Taxa <input type="text"/> Total <input type="text"/>	Comments: _____	

		Total Taxa
	<input type="text"/>	<input type="text"/>

Describe other aquatic life (e.g., fish, amphibians) collected or observed, as well as other indications that the reach is being used by other animals (i.e., birds, mammals, reptiles).

LEVEL-TWO SURVEY DATA SHEET

BIOLOGICAL INTEGRITY

The **SHADED** boxes indicate that multiple **FAMILIES** are possible; tolerance values are provided.

TV	Macroinvertebrates	Totals	Tolerance score	Number of kinds	TV	Macroinvertebrates	Totals	Tolerance score	Number of kinds	
1	Patterned stoneflies				6	Aquatic moth				
2	Winter stoneflies				4	Riffle beetle				
1	Roach-like stonefly				5	Long-toed beetle				
1	Giant stonefly				3	Water penny				
2	Little brown stonefly				5	Whirligig beetle				
3	Spiny crawler mayfly				7	Other beetles/bugs				
5	Square-gilled mayflies				3	Hellgrammite/Fishfly				
4	Minnow mayflies				6	Alderfly				
3	Flatheaded mayfly				9	Non-biting midge				
3	Brush-legged mayfly				6	Black fly				
5	Burrowing mayflies				4	Crane fly				
4	Net-spinning caddisflies				3	Watersnipe fly				
3	Case-building caddisflies				6	Dance fly				
5	Common netspinner				5	Dixid midge				
3	Free-living caddisfly				2	Net-wing midge				
4	Dragonflies				7	Horse fly				
7	Damselflies				8	Other fly larva				
Non-Insect Groups										
5	Crayfish				5	Pea clam				
5	Scud/Sideswimmer				6	Asian clam				
7	Aquatic sowbug				4	Mussel				
6	Water mite				5	Operculate snails				
10	Aquatic worms				7	Non-operculate snails				
10	Leeches				Other invertebrates					
7	Flatworms									
Complete your calculations using the metrics below. These metrics are combined to determine your overall score and integrity rating.	Total Number	Total Tolerance	Total Kinds	Comments: _____ _____ _____						

BSVs	Metrics	Results	Points	10	8	6	4	2
21	Total Taxa			> 20	19 - 16	15 - 12	11 - 8	< 8
14	EPT Taxa			> 12	12 - 9	8 - 5	4 - 2	< 2
3.20	Biotic Index			< 3.2	3.2 - 4.0	4.1 - 5.2	5.3 - 6.0	> 6.0
90.0	% EPT Abundance			> 85	85 - 75	74.9 - 65	64.9 - 50	< 50
81.4	% Dominance			< 10	10 - 15	15.1 - 25	25.1 - 50	> 50
2.0	% Tolerant			< 2	2 - 10	10.1 - 15	15.1 - 20	> 20
Stream Score				Integrity Rating				
				> 50	50 - 40	39 - 25	< 25	
				Optimal	Suboptimal	Marginal	Poor	

Another way to evaluate the benthic community is to use best standard values (BSVs). BSVs are used to calculate an overall score and integrity rating based on a 0-100 scale. [CLICK-HERE](#) to learn more.

Submit an original or clear copy of your survey to the [Coordinator](#) at the address provided below. For more information visit: <https://go.wv.gov/sos>

Philippi Field Office

Coordinator

WV Dept. of Environmental Protection
401 School Street, Suite 301
Philippi, WV 26416

Email: Callie.C.Sams@wv.gov
Office: (304) 314-6095
Mobile: (681) 341-9305

Attn: Callie Cronin Sams